

# An Improved Bound on the Number of Unit Area Triangles\*

Roel Apfelbaum<sup>†</sup>      Micha Sharir<sup>‡</sup>

January 26, 2010

## Abstract

We show that the number of unit-area triangles determined by a set of  $n$  points in the plane is  $O(n^{9/4+\varepsilon})$ , for any  $\varepsilon > 0$ , improving the recent bound  $O(n^{44/19})$  of Dumitrescu et al.

## 1 Introduction

In 1967, A. Oppenheim (see [5]) asked the following question: Given  $n$  points in the plane and  $A > 0$ , how many triangles spanned by the points can have area  $A$ ? By applying a scaling transformation, one may assume  $A = 1$  and count the triangles of *unit* area. Erdős and Purdy [4] showed that a  $\sqrt{\log n} \times (n/\sqrt{\log n})$  section of the integer lattice determines  $\Omega(n^2 \log \log n)$  triangles of the same area. They also showed that the maximum number of such triangles is at most  $O(n^{5/2})$ . In 1992, Pach and Sharir [7] improved the bound to  $O(n^{7/3})$ , using the Szemerédi-Trotter theorem [10] on the number of point-line incidences. Recently, Dumitrescu et al. [3] have further improved the upper bound to  $O(n^{44/19}) = O(n^{2.3158})$ , by estimating the number of incidences between the given points and a 4-parameter family of quadratic curves.

In this paper we further improve the bound to  $O(n^{9/4+\varepsilon})$ , for any  $\varepsilon > 0$ . Our proof borrows some ideas from [3], but works them into a different approach, which reduces the problem to bounding the number of incidences between points and certain kind of surfaces in three dimensions.

---

\*Work on this paper was supported by NSF Grants CCF-05-14079 and CCF-08-30272, by a grant from the U.S.-Israeli Binational Science Foundation, by grant 155/05 from the Israel Science Fund and by the Hermann Minkowski–MINERVA Center for Geometry at Tel Aviv University.

<sup>†</sup>School of Computer Science, Tel Aviv University, Tel Aviv 69978, Israel; roel6@hotmail.com.

<sup>‡</sup>School of Computer Science, Tel Aviv University, Tel Aviv 69978, Israel; and Courant Institute of Mathematical Sciences, New York University, New York, NY 10012, USA. E-mail: michas@post.tau.ac.il

## 2 Unit-area triangles in the plane

To simplify the notation, we write  $O^*(f(n))$  for an upper bound of the form  $C_\varepsilon f(n) \cdot n^\varepsilon$ , which holds for any  $\varepsilon > 0$ , where the constant of proportionality  $C_\varepsilon$  depends on  $\varepsilon$ .

**Theorem 2.1.** *The number of unit-area triangles spanned by  $n$  points in the plane is  $O^*(n^{9/4})$ .*

**Proof.** We begin by borrowing some notation and preliminary ideas from [3]. Let  $S$  be the given set of  $n$  points in the plane. Consider a triangle  $\Delta = \Delta abc$  spanned by  $S$ . We call the three lines containing the three sides of  $\Delta abc$ , *base lines* of  $\Delta$ , and the three lines parallel to the base lines and incident to the respective third vertices, *top lines* of  $\Delta$ .

For a parameter  $k$ ,  $1 \leq k \leq \sqrt{n}$ , to be optimized later, call a line  $\ell$  *k-rich* (resp., *k-poor*) if  $\ell$  contains at least  $k$  (resp., fewer than  $k$ ) points of  $S$ . Call a triangle  $\Delta abc$  *k-rich* if each of its three top lines is *k-rich*; otherwise  $\Delta$  is *k-poor*.

We first observe that the number of *k-poor* unit-area triangles spanned by  $S$  is  $O(n^2 k)$ . Indeed, assign a *k-poor* unit-area triangle  $\Delta abc$  whose top line through  $c$ , say, is *k-poor* to the opposite base  $ab$ . Then all the triangles assigned to a base  $ab$  are such that their third vertex lies on one of the two lines parallel to  $ab$  at distance  $2/|ab|$ , where that line contains fewer than  $k$  points of  $S$ . Hence, a base  $ab$  can be assigned at most  $2k$  triangles, and the bound follows.

So far, the analysis follows that of [3]. We now focus the analysis on the set of *k-rich* unit-area triangles spanned by  $S$ , and use a different approach.

Let  $L$  denote the set of *k-rich* lines, and let  $Q$  denote the set of all pairs

$$\{(\ell, p) \mid \ell \in L, p \in S \cap \ell\}.$$

By the Szemerédi-Trotter theorem [10], we have, for any  $k \leq \sqrt{n}$ ,  $m := |L| = O(n^2/k^3)$ , and  $N := |Q| = O(n^2/k^2)$ .

A pair  $(\ell_1, p_1), (\ell_2, p_2)$  of elements of  $Q$  is said to *match* if the triangle with vertices  $p_1, p_2, \ell_1 \cap \ell_2$  has area 1; see Figure 1.

To upper bound the number of unit-area triangles, all of whose three top lines are *k-rich*, it suffices to bound the number of matching pairs in  $Q$ . Indeed, given such a unit-area triangle  $\Delta p_1 p_2 q$ , let  $\ell_1$  (resp.,  $\ell_2$ ) be the top line of  $\Delta p_1 p_2 q$  through  $p_1$  (resp., through  $p_2$ ). Then  $(\ell_1, p_1)$  and  $(\ell_2, p_2)$  form a matching pair in  $Q$ , by definition (again, see Figure 1). Conversely, a matching pair  $(\ell_1, p_1), (\ell_2, p_2)$  determines at most one unit-area triangle  $p_1 p_2 q$ , where  $q$  is the intersection point of the line through  $p_1$  parallel to  $\ell_2$  and the line through  $p_2$  parallel to  $\ell_1$ ; we get an actual triangle if and only if the point  $q$  belongs to  $S$ .

In other words, our problem is now reduced to that of bounding the number of matching pairs in  $Q$ . (Since we do not enforce the condition that the third point  $q$  of the corresponding triangle belongs to  $S$ , we most likely over-estimate the true bound.)

Since elements of  $Q$  have three degrees of freedom, we can represent them in an appropriate 3-dimensional parametric space. For example, we can assume that no line in  $L$




Figure 1: The ordered pair  $((\ell_1, p_1), (\ell_2, p_2))$  is a matching pair of elements of  $Q$ .

is vertical, and parametrize an element  $(\ell, p)$  of  $Q$  by the triple  $(a, b, \kappa)$ , where  $(a, b)$  are the coordinates of  $p$ , and  $\kappa$  is the slope of  $\ell$ . For simplicity of notation, we refer to this 3-dimensional parametric space as  $\mathbb{R}^3$ .

So far, the matching relationship is symmetric. To simplify the analysis, and with no loss of generality, we make it asymmetric, by requiring that, in an (ordered) matching pair  $(\ell_1, p_1), (\ell_2, p_2)$ ,  $\vec{o}p_2$  lies counterclockwise to  $\vec{o}p_1$ , where  $o = \ell_1 \cap \ell_2$ . See Figure 1.

Let us express the matching condition algebraically. Let  $(a, b, \kappa) \in \mathbb{R}^3$  be the triple representing a pair  $(\ell, p)$ , and  $(x, y, w) \in \mathbb{R}^3$  be the triple representing another pair  $(\ell', p')$ . Clearly,  $w \neq \kappa$  in a matching pair. The lines  $\ell$  and  $\ell'$  intersect at a point  $o$ , for which there exist real parameters  $t, s$  which satisfy

$$o = (a + t, b + \kappa t) = (x + s, y + ws),$$

or

$$\begin{aligned} t &= \frac{y - b - w(x - a)}{\kappa - w} \\ s &= \frac{y - b - \kappa(x - a)}{\kappa - w}. \end{aligned}$$

It is now easy to verify that the condition of matching, with  $\vec{o}p'$  lying counterclockwise to  $\vec{o}p$ , is given by

$$\left( y - b - \kappa(x - a) \right) \left( y - b - w(x - a) \right) = 2(w - \kappa) \quad \text{and} \quad w \neq \kappa,$$

or, alternatively,

$$w = \frac{\left(y - b - \kappa(x - a)\right)(y - b) + 2\kappa}{\left(y - b - \kappa(x - a)\right)(x - a) + 2} \quad \text{and} \quad w \neq \kappa. \quad (1)$$

Similarly, the condition of “reverse” matching, with  $\vec{op}'$  lying clockwise to  $\vec{op}$ , is given by

$$w = \frac{\left(y - b - \kappa(x - a)\right)(y - b) - 2\kappa}{\left(y - b - \kappa(x - a)\right)(x - a) - 2} \quad \text{and} \quad w \neq \kappa. \quad (2)$$

Fix an element  $(\ell, p)$  of  $Q$ , and associate with it a surface  $\sigma_{\ell, p} \subset \mathbb{R}^3$ , which is the locus of all pairs  $(\ell', p')$  that match  $(\ell, p)$  (i.e.,  $(\ell, p), (\ell', p')$  is an ordered matching pair). By the preceding analysis,  $\sigma_{\ell, p}$  satisfies (1), where  $(a, b, \kappa)$  is the parametrization of  $(\ell, p)$ , and is thus a 2-dimensional algebraic surface in  $\mathbb{R}^3$  of degree 3. We thus obtain a system  $\Sigma$  of  $N$  2-dimensional algebraic surfaces in  $\mathbb{R}^3$ , and a set  $Q$  of  $N$  points in  $\mathbb{R}^3$ , and our goal is to bound the number of incidences between  $Q$  and  $\Sigma$ .

The main technical step in the analysis is to rule out the possible existence of *degeneracies* in the incidence structure, where many points are incident to many surfaces; this might happen when many points lie on a common intersection curve of many surfaces (a situation which might arise, e.g., in the case of planes and points in  $\mathbb{R}^3$ ). However, for the class of surfaces under consideration, namely, the surfaces  $\sigma_{\ell, p}$  generated by some line-point incidence pair  $(\ell, p)$ , such a degeneracy is impossible, as the following lemma shows.

**Lemma 2.2.** *Let  $(\ell_1, p_1)$  and  $(\ell_2, p_2)$  be two distinct line-point incidence pairs, let  $\gamma = \sigma_{\ell_1, p_1} \cap \sigma_{\ell_2, p_2}$  be the intersection curve of their associated surfaces, and assume that  $\gamma$  is non-empty. Let  $(\ell, p)$  be some incidence pair and assume further that  $\sigma_{\ell, p} \supset \gamma$ . Then either  $(\ell, p) = (\ell_1, p_1)$  or  $(\ell, p) = (\ell_2, p_2)$ .*

*Proof.* We establish the equivalent claim that, given a curve  $\gamma$ , which is the intersection of some unknown pair of surfaces  $\sigma_{\ell_1, p_1}$  and  $\sigma_{\ell_2, p_2}$ , one can reconstruct  $(\ell_1, p_1)$  and  $(\ell_2, p_2)$  uniquely (up to a swap between the two incidence pairs) from  $\gamma$ . Moreover, it is enough to know the projection  $\gamma^*$  of  $\gamma$  onto the  $xy$ -plane in order to uniquely reconstruct the incidence pairs  $(\ell_1, p_1)$  and  $(\ell_2, p_2)$  that generate  $\gamma$ .

We start by computing the algebraic representation of  $\gamma^*$ . Let  $(a_1, b_1, \kappa_1)$  and  $(a_2, b_2, \kappa_2)$  be the respective parametrizations of  $(\ell_1, p_1)$  and  $(\ell_2, p_2)$ . By (1),  $\gamma^*$  satisfies the equation

$$\frac{\left(y - b_1 - \kappa_1(x - a_1)\right)(y - b_1) + 2\kappa_1}{\left(y - b_1 - \kappa_1(x - a_1)\right)(x - a_1) + 2} = \frac{\left(y - b_2 - \kappa_2(x - a_2)\right)(y - b_2) + 2\kappa_2}{\left(y - b_2 - \kappa_2(x - a_2)\right)(x - a_2) + 2}. \quad (3)$$

Recall the additional requirement in (1), namely that  $w \neq \kappa_1$  and  $w \neq \kappa_2$ . This requirement is implicit in (1) and in (3), meaning that equation (3) is defined only for values of  $x$  and  $y$  for which the value of  $w$  is not  $\kappa_1$  or  $\kappa_2$ . Consulting (1), this implies that no point  $(x, y) \in \gamma$  can satisfy  $y - b_1 = \kappa_1(x - a_1)$  or  $y - b_2 = \kappa_2(x - a_2)$ . Put

$$\begin{aligned} L_1 &= y - b_1 - \kappa_1(x - a_1), & \text{and} \\ L_2 &= y - b_2 - \kappa_2(x - a_2), \end{aligned}$$

and write (3) as

$$\frac{L_1(y - b_1) + 2\kappa_1}{L_1(x - a_1) + 2} = \frac{L_2(y - b_2) + 2\kappa_2}{L_2(x - a_2) + 2},$$

or

$$(L_1(y - b_1) + 2\kappa_1)(L_2(x - a_2) + 2) = (L_2(y - b_2) + 2\kappa_2)(L_1(x - a_1) + 2),$$

which we can rewrite as

$$L_1L_2L_3 + 2L_1L_4 - 2L_2L_5 + 4C = 0,$$

where

$$\begin{aligned} L_3 &= (b_2 - b_1)x - (a_2 - a_1)y + (a_2b_1 - a_1b_2), \\ L_4 &= y - b_1 - \kappa_2(x - a_1), \\ L_5 &= y - b_2 - \kappa_1(x - a_2), \\ C &= \kappa_1 - \kappa_2. \end{aligned}$$

We can further simplify the equation by noting that  $L_6 = L_1L_4 - L_2L_5$  is a linear expression in  $x, y$ . That is,

$$L_6 = Dx + Ey + F,$$

where

$$\begin{aligned} D &= 2\kappa_1\kappa_2(a_2 - a_1) - (\kappa_1 + \kappa_2)(b_2 - b_1), \\ E &= 2(b_2 - b_1) - (\kappa_1 + \kappa_2)(a_2 - a_1), \\ F &= \kappa_1\kappa_2(a_1^2 - a_2^2) + (\kappa_1 + \kappa_2)(a_2b_2 - a_1b_1) + (b_1^2 - b_2^2). \end{aligned}$$

We can thus write (3) as

$$L_1L_2L_3 + 2L_6 + 4C = 0, \quad \text{and} \quad L_1 \neq 0, L_2 \neq 0. \quad (4)$$

Figure 2 illustrates the different lines defined by the linear equations  $L_i = 0$ , and their relations with  $(\ell_1, p_1)$  and  $(\ell_2, p_2)$ . The linearity of  $L_1, L_2, L_3$ , and  $L_6$  implies that the equation (4) of  $\gamma^*$  is cubic. We have the following two special cases to rule out:

1. If  $p_1 = p_2$ , that is,  $a_1 = a_2$  and  $b_1 = b_2$ , then  $L_3 = 0$ ,  $L_4 = L_2$ , and  $L_5 = L_1$ . But then the equation becomes  $4C = 0$ , or  $\kappa_1 = \kappa_2$ , contrary to the assumption that  $(\ell_1, p_1) \neq (\ell_2, p_2)$ . Hence, the equation has no solutions, meaning that  $\gamma$  is empty and the surfaces do not intersect.




Figure 2: The lines  $L_i = 0$ , for  $i = 1, \dots, 6$ , in the general case. The line  $L_3 = 0$  connects  $p_1$  and  $p_2$ ,  $L_4 = 0$  passes through  $p_1$  and is parallel to  $\ell_2$ ,  $L_5 = 0$  passes through  $p_2$  and is parallel to  $\ell_1$ , and  $L_6 = 0$  connects  $o = \ell_1 \cap \ell_2$  with the intersection point  $q$  of  $L_4 = 0$  and  $L_5 = 0$  (and bisects the edge  $p_1p_2$ ).

2. If  $\ell_1 = \ell_2$  but  $p_1 \neq p_2$ , that is,  $\kappa_1 = \kappa_2 = (b_2 - b_1)/(a_2 - a_1)$ , then  $L_1 = L_2 = L_4 = L_5$ ,  $L_3 = (a_1 - a_2)L_1$ , and  $C = 0$ , resulting in the equation  $(L_1)^3 = 0$ , which is not allowed in (4). Hence  $\gamma$  is not defined in this case either.

We can therefore restrict our attention to the general case. Consider the cubic part of the equation  $L_1L_2L_3$ . In this term, each factor can be thought of as a line defined by the equation  $L_i = 0$ , for  $i = 1, 2, 3$ . The lines  $L_1 = 0$  and  $L_2 = 0$  respectively are simply  $\ell_1$  and  $\ell_2$ , whereas  $L_3 = 0$  is the line  $\lambda$  passing through  $p_1$  and  $p_2$  (see Figure 2). Note that  $\lambda$  may coincide with one of the other two lines. Indeed, if  $p_1$  happens to be incident with  $\ell_2$ , then  $\lambda$  coincides with  $\ell_2$ . Similarly, if  $p_2 \in \ell_1$  then  $\lambda$  coincides with  $\ell_1$  (these are the only possible coincidences, since we have ruled out the case  $\ell_1 = \ell_2$ ). These cases will be handled shortly, but for now, we ignore them and consider the general case. In this case,  $\gamma^*$  has three distinct asymptotes given by  $L_1 = 0$ ,  $L_2 = 0$ , and  $L_3 = 0$ ; the proof of this fact is given in Lemma A.3 in the appendix

Using this fact, one can reconstruct the two line-point pairs that generate  $\gamma^*$  as follows. Suppose we are given a curve  $\gamma^*$  generated by some unknown pair of incidence pairs,  $(\ell_1, p_1)$  and  $(\ell_2, p_2)$ , and we want to reconstruct these pairs.  $\gamma^*$  is given as the zero set of some cubic bivariate polynomial  $f(x, y) = 0$ , where  $f$  can be written as  $f(x, y) = c(L_1L_2L_3 + 2L_6 + 4C)$ , but the decomposition of  $f$  into  $L_1, L_2, L_3, L_6, C$ , and  $c$  is unknown, and, moreover, is not known a priori to be unique (a fact which we establish in this proof). First, we find its three asymptotes  $\Lambda_1 = 0$ ,  $\Lambda_2 = 0$ , and  $\Lambda_3 = 0$ , where for each  $i = 1, 2, 3$ ,  $\Lambda_i$  is linear in  $x$  and  $y$ . Since, by Lemma A.3, these asymptotes are  $L_1 = 0$ ,  $L_2 = 0$ , and  $L_3 = 0$ , we know that each  $\Lambda_i$  is equal to some  $L_j$  multiplied by a constant, but we do not know which is which. To determine the roles of the asymptotes correctly, observe that  $\Lambda_1\Lambda_2\Lambda_3 = \mu L_1L_2L_3$  for some constant  $\mu$ . Thus, there exists some unique constant  $\nu$ , such that  $f(x, y) - \nu\Lambda_1\Lambda_2\Lambda_3 = \Lambda_4$  is linear in  $x$  and  $y$ . The line  $\Lambda_4 = 0$  is parallel to the line  $L_6 = 0$ , which happens to be the median of the triangle spanned by the three asymptotes, which emanates from the vertex  $o = \ell_1 \cap \ell_2$ , and bisects the edge  $p_1p_2$ ; see Figure 2. We thus have enough information to determine which vertex of the triangle is  $o$ , and which are  $p_1$  and  $p_2$ , and which edges of the

triangle are supported by  $\ell_1$  and  $\ell_2$ . This proves the lemma for the general case where all the points and lines are distinct, and no point lies on both lines  $\ell_2, \ell_2$ .

Finally, consider the case where  $p_2 \in \ell_1$  (a symmetric argument applies when  $p_1 \in \ell_2$ ). In this case,  $L_1 = L_5$ , and  $L_3 = (a_1 - a_2)L_1$ , so the equation of the curve  $\gamma^*$  can be rewritten as

$$(a_1 - a_2)L_1^2L_2 + 2L_1(L_4 - L_2) + 4C = 0.$$

Note that  $a_1 \neq a_2$  under the preliminary assumption that there are no vertical lines in the system, since both  $p_1 = (a_1, b_1)$  and  $p_2 = (a_2, b_2)$  are on  $\ell_1$ . Note also that  $C = \kappa_1 - \kappa_2 \neq 0$ , for otherwise,  $\ell_1$  and  $\ell_2$  would have to coincide, a case which we have ruled out earlier. Finally, note that  $s = L_4 - L_2 = b_2 - b_1 - \kappa_2(a_2 - a_1) = C(a_2 - a_1)$  is a nonzero constant. Hence, the equation of  $\gamma^*$  is, up to a constant multiple,

$$(a_1 - a_2)L_1^2L_2 + 2sL_1 + 4C = 0. \quad (5)$$

This equation defines a cubic curve with two asymptotes given by  $L_1 = 0$ , and  $L_2 = 0$ , namely, the lines  $\ell_1$  and  $\ell_2$ ; the proof is given in Lemma A.4 in the appendix. Since  $C \neq 0$ , it follows that  $\gamma^*$  does not intersect  $L_1 = 0$ , whereas  $L_2 = 0$  is intersected at a single point  $(x, y)$  for which  $L_1 = 2/(a_1 - a_2)$ . Using this point, one can compute the values of  $(a_1 - a_2)$ ,  $C$ , and  $s$ , and hence, reconstruct the line  $L_4 = 0$ . The point  $p_1$  is then simply the intersection of the lines  $L_1 = 0$  and  $L_4 = 0$ . Thus, one can uniquely reconstruct  $\ell_1$ ,  $\ell_2$ ,  $p_1$ , and  $p_2$  in this case too. This completes the proof of Lemma 2.2.  $\square$

**Bounding the number of incidences.** Recall that we need to bound the number of incidences between the set  $\Sigma$  of surfaces  $\sigma_{\ell, p}$ , for  $(\ell, p) \in Q$ , and the set  $Q$  of points. This is done by following the standard method of Clarkson et al. [2]. The first step in this method is to derive a simple but weaker bound, usually by extremal graph theory. Then, we strengthen the bound by *cutting* the arrangement of the surfaces into cells, and by summing the weaker bounds on the number of incidences within each cell, over all the cells.

**The first step: A simple bound.** Lemma 2.2 implies that the incidence graph between  $\Sigma$  and  $Q$  does not contain  $K_{3,10}$  as a subgraph, or, in other words, no three distinct surfaces of  $\Sigma$  and ten distinct points of  $Q$  can all be incident to one another. Indeed, the intersection points of three surfaces  $\sigma_{\ell_i, p_i}$ , for  $i = 1, 2, 3$ , are the intersection points of the two curves  $\gamma_{1,2} = \sigma_{\ell_1, p_1} \cap \sigma_{\ell_2, p_2}$ , and  $\gamma_{1,3} = \sigma_{\ell_1, p_1} \cap \sigma_{\ell_3, p_3}$ . These intersection points project to (some of) the intersection points of the projections  $\gamma_{1,2}^*$  and  $\gamma_{1,3}^*$  of  $\gamma_{1,2}$  and  $\gamma_{1,3}$ , respectively, onto the  $xy$ -plane. By Lemma 2.2, these two curves are distinct (or empty). Since each of them is cubic, and since, as shown in Lemmas A.3 and A.4 in the appendix, they are the zero sets of irreducible polynomials, Bézout's theorem [8] implies that they intersect in at most  $3^2 = 9$  points. Hence, the incidence graph between  $\Sigma$  and  $Q$  does not contain  $K_{3,10}$ , so by the Kővari–Sós–Turán theorem [6], the number of incidences between  $\Sigma$  and  $Q$  can be bounded by

$$O(|\Sigma||Q|^{2/3} + |Q|).$$

Since the matching relation is essentially symmetric (up to some sign changes; see (1) and (2)), we can interchange the roles of points and surfaces, and conclude that the number of incidences is also at most

$$O(|Q||\Sigma|^{2/3} + |\Sigma|). \quad (6)$$

**Cutting.** To improve the bound, we apply the following fairly standard space decomposition technique. Fix a parameter  $r$ , whose specific value will be chosen later, and construct a  $(1/r)$ -cutting  $\Xi$  of  $\mathcal{A}(\Sigma)$  [1]. We use the more simple-minded technique in which we choose a random sample  $R$  of  $O(r \log r)$  surfaces of  $\Sigma$  and construct the vertical decomposition (see e.g. [9]) of the arrangement  $\mathcal{A}(R)$ . We obtain  $O^*(r^3)$  relatively open cells of dimensions 0, 1, 2, and 3, each of which is crossed by (intersected by, but not contained in) at most  $|\Sigma|/r = N/r$  surfaces; this latter property holds with high probability, and we simply assume that our sample  $R$  does satisfy it.

**Summing over all cells.** Fix a cell  $\tau$  of  $\Xi$ , and put  $Q_\tau := Q \cap \tau$  and  $m_\tau := |Q_\tau|$ . Let  $\Sigma_\tau$  denote the subset of surfaces of  $\Sigma$  which cross  $\tau$ , and put  $N_\tau := |\Sigma_\tau| \leq N/r$ .

We now apply the simple bound (6) obtained in the first step to each cell  $\tau$  of our cutting  $\Xi$ , handling, for the time being, only surfaces that *cross*  $\tau$ . The overall number of incidences is

$$\sum_{\tau \in \Xi} O(m_\tau N_\tau^{2/3} + N_\tau),$$

which, using the bounds  $N_\tau \leq N/r$ , and  $\sum_\tau m_\tau = N$ , is

$$O^*(N(N/r)^{2/3} + Nr^2) = O^*(N^{5/3}/r^{2/3} + Nr^2).$$

To minimize this expression, we choose  $r = N^{1/4}$ , making it  $O^*(N^{3/2})$ .

We also have to take into account incidences between points in a cell  $\tau$  and surfaces that fully contain  $\tau$ . This is done separately for cells of dimension 0, 1, and 2 (it is vacuous for cells of dimension 3). Indeed, a 2-dimensional cell  $\tau$  is contained in exactly one surface, so a point  $w \in \tau$  takes part in only one such incidence. Thus, in this case we only need to add  $N$ , the number of points, to the above bound.

The same argument applies for points in 1-dimensional cells. Assuming that the vertical decomposition is performed in a generic coordinate frame, it suffices to consider only 1-dimensional cells that are portions of the intersection curves between the surfaces of  $\Sigma$ . By Lemma 2.2, each such cell  $\tau$  is contained in exactly two surfaces of  $\Sigma$ . Thus, we need to add at most  $2N$  to the number of incidences to handle these cells.

Each cell of dimension 0 is a single point  $w$ , and, arguing as above, we may assume it to be a vertex of the undecomposed arrangement  $\mathcal{A}(R)$ . Any surface  $\sigma$  incident to  $w$  has to cross or bound an adjacent full-dimensional cell  $\tau^*$ , so we charge the incidence of  $\sigma$  with  $w$  to the pair  $(\tau^*, \sigma)$ , and note that such a pair can be charged only  $O(1)$  times. It follows that the number of incidences with 0-dimensional cells of  $\Xi$  is  $O^*(r^3 + r^3(N/r)) = O^*(r^2N)$ , which, for the chosen value of  $r$ , is equal to the bound obtained above for the crossing surfaces.



In conclusion, the overall number of incidences between  $\Sigma$  and  $Q$  is  $O^*(N^{3/2})$ .

Recall now that  $N = O(n^2/k^2)$ , and that we also have the bound  $O(n^2k)$  for the number of unit-area triangles with at least one  $k$ -poor top line. Thus, the overall bound on the number of unit-area triangles is

$$O^*\left(\frac{n^3}{k^3} + n^2k\right),$$

which, if we choose  $k = n^{1/4}$ , becomes  $O^*(n^{9/4})$ , as asserted.  $\square$

**Discussion.** Theorem 2.1 constitutes a major improvement over previous bounds, but it still leaves a substantial gap from the near-quadratic lower bound. One major weakness of our proof is that, in bounding the number of matching pairs, it ignores the constraint that a matching pair is relevant only when the (uniquely defined) third vertex  $q$  of the resulting triangle belongs to  $S$ , and that the (uniquely defined) top line of this triangle through  $q$  is  $k$ -rich. It is therefore natural to conjecture that our bound is not tight, and that the true bound is nearly quadratic, perhaps coinciding with the lower bound of [4].

## References

- [1] B. Chazelle, Cuttings, In *Handbook of Data Structures and Applications* (D. Mehta and S. Sahni, editors), chap. 25, Chapman and Hall/CRC Press, 2005.
- [2] K. Clarkson, H. Edelsbrunner, L. Guibas, M. Sharir and E. Welzl, Combinatorial complexity bounds for arrangements of curves and spheres, *Discrete Comput. Geom.* 5 (1990), 99–160.
- [3] A. Dumitrescu, M. Sharir and Cs. D. Tóth, Extremal problems on triangle areas in two and three dimensions, *J. Combinat. Theory, Ser. A*, accepted. Also in in *Proc. 24th ACM Symp. on Computational Geometry* (2008), 208–217.
- [4] P. Erdős and G. Purdy, Some extremal problems in geometry, *J. Combinat. Theory* 10 (1971), 246–252.
- [5] P. Erdős and G. Purdy, Extremal problems in combinatorial geometry. in *Handbook of Combinatorics* (R. Graham, M. Grötschel and L. Lovász, editors), Vol. 1, 809–874, Elsevier, Amsterdam, 1995.
- [6] T. Kővari, V. T. Sós, and P. Turán, On a problem of K. Zarankiewicz, *Colloquium Math.* 3 (1954), 50–57.
- [7] J. Pach and M. Sharir, Repeated angles in the plane and related problems, *J. Combinat. Theory Ser. A* 59 (1992), 12–22.
- [8] I. R. Shafarevich, *Basic Algebraic Geometry*, Springer-Verlag, 1977.

- [9] M. Sharir and P. K. Agarwal, *Davenport-Schinzel Sequences and Their Geometric Applications*, Cambridge University Press, New York, 1995.
- [10] E. Szemerédi and W. T. Trotter, Extremal problems in discrete geometry, *Combinatorica* 3 (1983), 381–392.

## A Asymptotes of cubic curves

In this appendix, we analyze the class of cubic curves defined by equations (4) and (5) of Section 2, derive their asymptotes, and show them to be the zero sets of irreducible bivariate polynomials. We start by analyzing a normalized version of these equations, in which two of the generating lines (and, as we show henceforth, the asymptotes) are the  $x$  and  $y$ -axes. We then reduce equation (4) to the normalized case. Finally, we handle equation (5) in a different and simpler way.

**Lemma A.1.** *Let  $\lambda_1$  and  $\lambda_2$  be two distinct lines in  $\mathbb{R}^2$ , given by the equations  $\Lambda_i = 0$ , where  $\Lambda_i = \alpha_i x + \beta_i y + \gamma_i$ , and  $\alpha_i$  and  $\beta_i$  are both nonzero, for  $i = 1, 2$ . Let  $f(x, y)$  be the bivariate cubic polynomial*

$$f(x, y) = xy\Lambda_1 + \Lambda_2.$$

*Then  $f(x, y)$  is irreducible.*

*Proof.* Assume, to the contrary, that  $f$  is reducible. Then it has a linear factor  $L = ax + by + c$ . Without loss of generality,  $b \neq 0$  (a symmetric argument follows for the case  $a \neq 0$ ), so we can assume  $b = 1$ . Then  $f$ , as a polynomial in  $y$  with coefficients from  $\mathbb{R}[x]$ , has  $y = -ax - c$  as root, i.e., if we put

$$p(x) := f(x, -ax - c) = -x(ax + c)L_1 + L_2,$$

where  $L_i = \alpha_i x - \beta_i(ax + c) + \gamma_i$ , for  $i = 1, 2$ , then  $p(x) \equiv 0$ . But then, the term  $x(ax + c)L_1$  can not be properly cubic, nor quadratic, so,  $a = c = 0$ , or  $L_1$  is a constant, possibly zero. In the former case,  $L_2 \equiv 0$ , but  $L_2 = \alpha_2 x + \gamma_2$  and  $\alpha_2 \neq 0$  by assumption, a contradiction. If  $L_1 = 0$ , then we must also have  $L_2 = 0$ , and so both lines  $\lambda_1$  and  $\lambda_2$  coincide (with the line  $L = 0$ ), contrary to assumption. If  $L_1$  is a nonzero constant, then the term  $(ax + c)$  must also be constant, or else  $p(x)$  is a proper quadratic polynomial, hence  $a = 0$ . But then, for  $L_1 = \alpha_1 x - \beta_1 c + \gamma_1$  to be constant, we must have  $\alpha_1 = 0$ , in contradiction. Either way,  $f$  cannot be reducible.  $\square$

**Lemma A.2.** *Let  $\lambda_1$  and  $\lambda_2$  be two distinct lines in  $\mathbb{R}^2$ , given by the equations  $\Lambda_i = 0$ , where  $\Lambda_i = \alpha_i x + \beta_i y + \gamma_i$ , for  $i = 1, 2$ , such that  $\alpha_1$  and  $\beta_1$  are both nonzero. Let  $\Gamma$  be the algebraic cubic curve defined by the equation*

$$xy\Lambda_1 + \Lambda_2 = 0. \tag{7}$$

*Then  $\Gamma$  is asymptotic to the  $x$ -axis and to the  $y$ -axis.*

*Proof.* We only prove in detail that the  $x$ -axis is an asymptote. Note that, for any fixed  $x \neq 0$ , (7) is a quadratic equation in  $y$ , which we rewrite as

$$xy(\alpha_1x + \beta_1y + \gamma_1) + \alpha_2x + \beta_2y + \gamma_2 = 0,$$

or

$$\beta_1xy^2 + (\alpha_1x^2 + \gamma_1x + \beta_2)y + (\alpha_2x + \gamma_2) = 0.$$

Hence

$$y = -\frac{\alpha_1x^2 + \gamma_1x + \beta_2}{2\beta_1x} \pm \frac{\sqrt{(\alpha_1x^2 + \gamma_1x + \beta_2)^2 - 4\beta_1x(\alpha_2x + \gamma_2)}}{2\beta_1x}.$$

We only consider the solution with positive square root, which is

$$y = \frac{\alpha_1x^2 + \gamma_1x + \beta_2}{2\beta_1x} \left[ \sqrt{1 - \frac{4\beta_1x(\alpha_2x + \gamma_2)}{(\alpha_1x^2 + \gamma_1x + \beta_2)^2}} - 1 \right].$$

The expression in the square brackets is of the form  $\sqrt{1+t} - 1$ . Since  $\alpha_1 \neq 0$ ,  $t$  tends to 0 as  $x \rightarrow \pm\infty$ . Using the inequalities  $1 - |t| \leq \sqrt{1+t} \leq 1 + \frac{|t|}{2}$ , for  $|t| < 1$ , we obtain, for  $|x|$  sufficiently large,

$$|y| \leq \frac{|\alpha_1x^2 + \gamma_1x + \beta_2|}{|2\beta_1x|} |t| = \frac{2|\alpha_2x + \gamma_2|}{|\alpha_1x^2 + \gamma_1x + \beta_2|},$$

which tends to 0 as  $x \rightarrow \pm\infty$ . This shows that the  $x$ -axis is indeed an asymptote of  $\Gamma$  (on both sides). A symmetric argument shows that the  $y$ -axis is also an asymptote.  $\square$

We are now ready to prove the more general cases discussed in Section 2.

**Lemma A.3.** *Let  $\ell_1, \dots, \ell_4$  be four distinct lines in  $\mathbb{R}^2$ , given by the equations  $L_i = 0$ , where  $L_i = A_ix + B_iy + C_i$ , for  $i = 1, \dots, 4$ . Assume that no pair of  $\ell_1, \ell_2, \ell_3$  are parallel, and that  $\ell_4$  is not parallel to any of  $\ell_1$  and  $\ell_2$ . Put*

$$f(x, y) = L_1L_2L_3 + L_4,$$

*and let  $\Gamma$  be the algebraic cubic curve defined by the equation*

$$f(x, y) = 0.$$

*Then,  $f$  is irreducible, and  $\Gamma$  is asymptotic to the lines  $\ell_1, \ell_2, \ell_3$ .*

*Proof.* We may assume, by an appropriate change of variables, that one of  $\ell_1, \ell_2$ , and  $\ell_3$  is the  $x$ -axis and another one is the  $y$ -axis. For example, put  $u = L_1$ , and  $v = L_2$ , and write  $L_3 = \alpha_1u + \beta_1v + \gamma_1$ , and  $L_4 = \alpha_2u + \beta_2v + \gamma_2$ , for some appropriate coefficients  $\alpha_1, \beta_1, \gamma_1, \alpha_2, \beta_2, \gamma_2$ . Note that, by the preliminary assumptions on the lines,  $\alpha_i$  and  $\beta_i$  are both nonzero, for  $i = 1, 2$ .  $\Gamma$  can then be written as

$$g(u, v) = uvL_3 + L_4 = 0$$

in the  $(u, v)$  coordinate system. It then follows, by Lemma A.1, that  $f$  is irreducible, for otherwise, any factorization of  $f$  could be transformed into a factorization of  $g$ , in contradiction. It also follows, by Lemma A.2, that  $\ell_1$  and  $\ell_2$  are asymptotes of  $\Gamma$ . Note that, for this part of the argument, the choice of  $\ell_1$  and  $\ell_2$  as axes is arbitrary, and we could just as well choose any other pair of lines among  $\ell_1, \ell_2, \ell_3$ . In more detail, since no pair of these three lines are parallel, we can make any two of them as the axes of a new  $(u, v)$ -coordinate system, and then, in the equation of the third line, both the  $u$ - and  $v$ -coefficients would be nonzero, which is the condition assumed in Lemma A.2. Hence,  $\ell_3$  is also an asymptote of  $\Gamma$ .  $\square$

**Lemma A.4.** *Let  $\ell_1$  and  $\ell_2$  be two distinct intersecting lines in  $\mathbb{R}^2$ , given by the equations  $L_i = 0$ , where  $L_i = A_i x + B_i y + C_i$ , for  $i = 1, 2$ . Put  $f(x, y) = L_1^2 L_2 + L_1 + C$ , for some constant  $C$ , and let  $\Gamma$  be the algebraic curve defined by the equation*

$$f(x, y) = 0.$$

*Then  $\Gamma$  is asymptotic to the lines  $\ell_1$  and  $\ell_2$ . Furthermore, if  $C \neq 0$ , then  $f$  is an irreducible bivariate polynomial.*

*Proof.* If  $C = 0$ , then the claim is easy. Indeed, in this case we have  $L_1(L_1 L_2 + 1) = 0$ , so  $\Gamma$  is the union of the line  $L_1 = 0$  and the hyperbola  $L_1 L_2 = -1$ , which is asymptotic to the lines  $L_1 = 0$ , and  $L_2 = 0$ .

If  $C \neq 0$ , put  $u = L_1$ , and  $v = L_2$ . Then, in the  $(u, v)$  coordinate system,  $\Gamma$  is defined by the equation

$$g(u, v) := u^2 v + u + C = 0.$$

Note that  $g$  is clearly irreducible, and so is  $f$ . This equation can be rewritten as

$$v = -\frac{u + C}{u^2}.$$

Clearly, this function tends to 0 as  $u$  tends to  $\infty$ , which means that  $\Gamma$  is asymptotic to the  $u$ -axis, i.e., to  $\ell_2$ . Furthermore, the function has a pole at  $u = 0$ , meaning that  $\Gamma$  is asymptotic to the  $v$ -axis, i.e., to  $\ell_1$ .  $\square$